



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

disturbances following the first attack, since just beneath this region the sensory fibers for the entire body are most compactly grouped. The autopsy fully confirmed the diagnosis. There were several small cortical lesions and an old lesion in the pons—all without significance for the present discussion. Besides these, however, there was a fresh lesion in the third member of the right lenticular nucleus, which would easily involve the pyramidal tract, while the white matter of the lower portion of the right parietal lobe was completely softened, thus fully accounting for the other symptoms. A review of the literature brought to light a considerable amount of evidence showing conjugate deviation of the eyes in similar lesions of the parietal lobe.

Die Beziehungen der hinteren Rindengebiete zum epileptischen Anfall.
H. UNVERRICHT. Deutsch. Archiv f. klin. Med. XLIV, 1, S. 1.
Reviewed by Ziehen, Centralbl. f. Physiol. No. 25, 1889.

The starting point for an epileptic attack is usually thought of as in the motor region of the cortex, but Unverricht considers it as clinically established that a strictly local affection of the posterior cortical regions can of itself bring about convulsions. He seeks to demonstrate the point on dogs, in which the stimulation, especially of the posterior and superior portions of the second arched convolution (counting from the middle line) causes contractions. There is lateral motion of the eyes to the opposite side, with dilation of the pupils as one of the results. These are not explained as reflexes from sensory stimuli (Ferrier), but as the result of direct stimulation of motor centers. But this is simply the author's view, for which the evidence is lacking thus far.

The length of time the stimulus is continued is more important than the strength of the stimulus, in determining an attack from this posterior cortical area. The order of contractions often fails to follow the order of the centers, and at times the convulsions roused from one visual area are limited to one half of the body. Most important are two experiments in which on the left side all the motor region save that for the movements of the eyes had been removed. The visual area was then stimulated on the left side, and the convulsion appeared on the *same* side. When a transverse cut was made through the cortex of one hemisphere at the anterior edge of the visual area, then stimulation of the latter caused a convulsion in which the orbicularis contracted after the extremities, from which he concludes that the impulse travels through deep-lying connections. At the end some clinical evidence is presented.

Kleine Beiträge, betreffend die Anordnung der Geschmacksknospen bei den Säugethieren. J. HÖNIGSCHMIED. Zeitschr. f. wiss. Zool., Vol. XLVII, 1888, S. 190-200.

This paper contains the results of the author's further studies upon the arrangement of the gustatory papillae and distribution of the taste-bulbs in mammals. In *Felis lynx* there are present six papillae of the circumvallate type, but the foliate papillae are wanting. The taste-bulbs are rather narrow and disposed in a zone of 3 to 5 tiers. *Ursus fuscus* has about twenty circumvallate papillae, and also well-developed foliate organs. The taste-bulbs are oval or cylindrical in form and, in the circumvallate papillae, are arranged